



Are you being treated for atrial fibrillation (Afib) but still experiencing these symptoms?

- Shortness of breath
- Weakness
- Fatigue
- Pain or pressure in chest
- Lowered blood pressure
- Dizziness
- Rapid or irregular heartbeat

Some people with Afib may not be aware of their symptoms and are only diagnosed through an assessment by their physician.

It may be time to talk to your provider about a change in your treatment plan.


Ask your physician today if you are a candidate for Hybrid AF Therapy.

Each patient's treatment plan is unique. But if Afib is not adequately treated, it may cause physical changes to your heart. It may also increase your risk for stroke, and heart failure.



Visit www.HybridAFTherapy.com for additional information and resources.

Schedule an appointment today with your Heart Team:



Hybrid AF Therapy is for the treatment of long-standing persistent atrial fibrillation.

Risk Information: This procedure is not recommended for patients with Barrett's Esophagitis, presence of left atrial thrombus (clot), a systemic infection, or an active infection local to the surgical site at the time of surgery (i.e. active endocarditis).

Potential procedural complications include, but are not limited to: Pericardial effusion, pericarditis, infection, cardiac tamponade, pulmonary vein stenosis, vessel injury, tissue perforation, excessive bleeding, phrenic nerve injury, left atrial rupture, esophageal fistula, heart attack, new arrhythmias, thromboembolic complication, stroke/TIA/neurologic complication, complete heart block requiring permanent pacemaker implantation, serious skin burn, a buildup of fluid around your lungs, or death.

This information is not comprehensive. Talk to your health care provider to obtain the FDA-approved product labeling or visit www.AtriCure.com.

Rx Only.

PM-US-1519C-0825-G